

INTERNATIONAL CENTRE FOR ADVANCED MEDITERRANEAN AGRONOMIC STUDIES MEDITERRANEAN AGRONOMIC INSTITUTE OF ZARAGOZA



Advanced Course

FUNCTIONAL FOODS: SCIENTIFIC FOUNDATIONS AND OPPORTUNITIES FOR THE AGRO-FOOD SECTOR

Zaragoza (Spain), 3-7 April 2006





CENTRO INTERNACIONAL DE ALTOS ESTUDIOS AGRONOMICOS MEDITERRANEOS INSTITUTO AGRONOMICO MEDITERRANEO DE ZARAGOZA



Curso Avanzado

ALIMENTOS FUNCIONALES: BASES CIENTÍFICAS Y OPORTUNIDADES PARA EL SECTOR AGROALIMENTARIO

Zaragoza (España), 3-7 abril 2006



Francisco Javier Señoráns

Dept. of Food Science
Universidad Autónoma de Madrid







FUNCTIONAL FOODS: SCIENTIFIC FOUNDATIONS AND OPPORTUNITIES FOR THE AGRO-FOOD SECTOR



Francisco Javier Señoráns

Dept. of Food Science
Universidad Autónoma de Madrid





FUNCTIONAL FOODS: SCIENTIFIC FOUNDATIONS AND OPPORTUNITIES FOR THE AGRO-FOOD SECTOR


Functional ingredients and their effects:

Lipophilic Ingredients from vegetable products



Dr. Francisco Javier Señoráns

Dept. of Food Science
Universidad Autónoma de Madrid (Spain)





FUNCTIONAL FOODS



Lipidic Ingredients from vegetable products

Functional Foods

Son alimentos que tienen un efecto beneficioso sobre una o más funciones en el cuerpo, por encima y más allá de los efectos habituales de dicho alimento, tales como mejorar el estado de salud y bienestar o reducir el riesgo de enfermedad.

European International Life Sciences Institute (LSI)











FUNCTIONAL FOODS



Lipidic Ingredients from vegetable products

Making of completely new Foods.....



















...or extraction of bioactive and health promoting ingredients from other products



FUNCTIONAL FOODS

Lipophilic Ingredients from vegetable products

Bioactive and health promoting ingredients from vegetable products



Potential functional food ingredients





Functional Ingredients from vegetable products

Bioactive and health promoting ingredients from vegetable products

- 1. FATTY ACIDS (OLEIC, LINOLENIC, OMEGA-3, ETC.)
- 2. TOCOPHEROLS
- 3. STEROLS
- 4. CAROTENOIDS
- 5. OTHER ANTIOXIDANTS: PHENOLIC DITERPENES
- 6. ISOFLAVONES AND ISOFLAVONOIDS
- 7. FLAVONOIDS
- 8. PHENOLIC COMPOUNDS
- 9. ORGANOSULFUR COMPOUNDS
- 10. ETC...



FUNCTIONAL FOODS

Lipophilic Ingredients from vegetable products

Which lipophilic plant ingredients?

- TOCOPHEROLS
- STEROLS
- CAROTENOIDS
- OTHER ANTIOXIDANTS: PHENOLIC DITERPENES



FUNCTIONAL FOODS

Lipophilic Ingredients from vegetable products

Which lipophilic plant ingredients?

- **TOCOPHEROLS**
- STEROLS
- CAROTENOIDS
- OTHER ANTIOXIDANTS: PHENOLIC DITERPENES.

Research and development on these ingredients?





Lipidic Ingredients from vegetable products

Tocopherols

Sterols Carotenoids

Other antioxidants

What are they? Structure - types-function

Why functional ingredients? (bioactivity)

Do you have scientific evidence? Articles

Do you have clinical studies? Health claims



OK, Lets do it

Where are they? Sources-industrial production

What functional food is the ideal for this ingredient? Available, compatible, complementary

Produce and sell it!



VITAMIN E...

...Functional food ingredient?

5.0

TOCOPHEROLS

Vitamin E

- The term vitamin E refers to a family of eight related compounds, the tocopherols and the tocotrienols
- The four major forms of vitamin E are designated α , β , δ , γ , on the basis of the number and position of the methyl group on the chromanol ring

Structure

•alpha-tocopherol

alpha-tocotrienol

TOCOPHEROLS

What are they? Structure - types-function



- Bioactivity of different Isomers: Vitamin E •4 tocopherols + 4 tocotrienols
- •alpha-tocopherol 3 chiral centers

- •Part of antioxidant network in biological systems (with Vitamin C)

•Synthetic is all-rac-alpha-tocopherol (mixture of 8 isomers)






Why functional ingredients? (bioactivity)


- •Biological effect: Vitamin E activity:
 - •RRR-alpha-tocopherol 1.49 IU /mg highest activity
- •Alpha-tocopherol ---highest bioavailability
 - predominant in human plasma and tissues
 - selective incorporation into nascent VLDL


TOCOPHEROLS

Bioactivity and function

- Antioxidant
- · Free radical scavenger
- Scavenge lipid peroxyl radicals
- Efficient oxygen quencher
 - Protects cell membranes
 - Protects LDL from oxidation
 - Prevention of rancidity
 - Works in conjunction with selenium
- Protection of double bonds in polyunsaturated fatty acids
 - Protects lungs from pollutants
 - Protects DNA
 - Protects heartMain activity of Tocopherols:
 - Others? Most important lipid soluble antioxidant









Is there any scientific evidence?

- •Chemical characterization and Different tests (activity *invitro...*): hundreds
- •Epidemiological studies: several evidences
- •Animal model tests (in-vivo)
- Pre-clinical: Dose effects, toxicological risks, etc.
- •Clinical studies, Intervention studies: many
- •Reliability: Total
- •Controversial:
 - None on effect
 - Possible on need of supplementation
- Accepted: yes
- Approved health claims:

•??

TOCOPHEROLS

Are there clinical studies? Health claims

- Biological activity
 - Antioxidant
- Health promoting effects:
 - Protection against Atherosclerosis and Coronary Heart Disease
 Reduce LDL oxidation: numerous studies (Lapointe, 2006)
 - Protection against some cancers: under study
 hormonal, digestive tract...
- **Recommended daily intake: 10 mg** α -tocopherol/day (15 IU)
- •Supplementation 15-30 mg a-tocopherol/day to 100mg
- •Related to intake of PUFA



Do you have clinical studies? Health claims

A. Lapointe et al. / Journal of Nutritional Biochemistry xx (2006) xxx-xxx

Table 1

THOSE I		
Studies investigating the effects of autoconhere	I supplementation on LDI.	ovidation parameters

Authors (date)	Subjects' characteristics	Study design/duration	Dosage (per day)	Outcomes
Reaven et al. (1993) [66]	Mildly hyperlipidemic subjects (n=16)	Randomized Free-living Supplement provided 8 weeks	1600 mg, control	Lag phase with AT after 4 and weeks compared to control Maximum rate of propagation
Jialal et al, (1995) [60]	Healthy subjects (n=48)	Randomized Free-living Supplements provided 8 weeks	60, 200, 400, 800 or 1200 IU, placebo (soybean oil)	↑ Lag phase with AT (400, 800 and 1200 IU/day) compared to baseline ↓ Oxidation rate (CD) with AT (800 and 1200 IU/day) compared to baseline
Princen et al. (1995) [59]	Healthy subjects (n=20)	Parallel Free-living Supplements provided 2 weeks	25, 50, 100, 200, 400 or 800 IU	↑ Lag phase with AT (25, 100, 200, 400 an 800 IU/day) compared to baseline ↓ Propagation rate with AT (400 and 800 UI/day) compared to previous dose
Reaven et al. (1995) [64]	T2D males (n = 21)	Randomized Free-living Supplements provided	1600 IU or placebo	↑ Lag phase with AT compared to previous dose
Suzukawa et al. (1995) [58]	Healthy subjects (n=8)	Free-living Supplements provided 4 weeks	150 mg (1 week) and 300 mg (3 weeks)	↑ Lag phase with AT compared to baseline ↓ Propagation rate compared to baseline
Fuller et al. (1996) [65]	Diabetic subjects (n=28)	Randomized Free-living Supplements provided 8 weeks	1200 IU or placebo	↑ Lag phase with AT compared to baseline and to placebo ↓ Oxidation rate (CD) with AT compared to baseline
Mol et al. (1997) [61]	T2D subjects (n=11) Healthy subjects (n=14)	Free-living Supplements provided	600 IU	↑ Lag phase in three groups compared to baseline

FUNCTIONAL INGREDIENTS



Lipidic Ingredients from vegetable products

Tocopherols

What are they? Structure - types-function

Why functional ingredients? (bioactivity)

Do you have scientific evidence? Articles

Do you have clinical studies? Health claims



OK, Lets make it

Where are they? Sources-industrial production

What functional food is the ideal for this ingredient? Available, compatible, complementary

Different cases:tocos-sterols-carots-etc



Where are they? Natural sources

- · Found in all photosynthetic organisms
 - •always located in membrane of the cell
- •Sources in diet:
 - Fats and oils (corn, soybean, palm, cottonseed, sunflower..)
 - Cereal grains (minor)
 - Vegetable products
- •Effect of growing conditions
 - · Higher levels at higher temperatures




TOCOPHEROLS

Where are they? Industrial production

- Extraction from natural sources
 - •Deodorization distillates from edible oil processing
 - •From soybean oil (10-14% tocos) or corn
- Chemical synthesis
 - •Synthesis of 8 stereoisomers- high yield---esterification (stable)—purification (vacuum distillation)



Where are they? Sources-industrial production



TOCOPHEROLS



Where are they? Sources-industrial production

Eur Food Ros Technol (2003) 217:427-413 pon 10:1007/st0217-403-0761-5

ORIGINAL PAPER

G. K. Nagesha - B. Manshar - K. Udaya Sankar

Enrichment of tocopherols in modified soy deodorizer distillate using supercritical carbon dioxide extraction

Rosson 17 February 2005 / Published selline: 1 October 2007

Admiratel suppositional carbon broads (20°C-32) establishment between the CDO3 and the electric (1905) at effect and of procure (1905) 400 stars and temperature (1956) Cr. to reptains the conditions for temperature (1956) Cr. to reptains the conditions for the condition of the

Keywords Soy DOD (decelerater distillate) - Medified soy DOD - Supercritical carbon distrilet (SC-CO₂) - Encolment - Tecopherols - Fatty acid methyl enters

Introduction

Sey decolarizer discillate (DOD) in a valuable by-produc in the refining of vegetable rils and amounts to 0.3-0.59 G. R. Sugenho - B. Manshar - K. Udeya Santar (20) Department of Fixed Engineering, Control Food Technological Research Institute. of the first cock, h is complex nations of subgrounds for first paids, term by explored, behaviors and suffer susquestful behaviors and other susquestful behaviors and enter susquestful behaviors. Among these two pleases in control statement agriculture in content, plantum control statement and on natural and entitles in proceeding the control of the surface of the surface and the first statement and such large statement and such large statement and surface activates of fast and sub-first attempts and the proceedings. Principality the entities in some law plantum designs after the surface of the surface and a plantum containers. Principality the entities in some statement and the surface and the surface and the surface of the surface and th

Secondary Companies of the Companies of

Supervisical carbon densitie (SC CO2) extraction received as a speciation go as speciation go to a special properties go as special probability of the probability of



Food Chemistry 85 (2004) 579-583

Food Chemistry

Developing functional foods using red palm olein. IV. Tocopherols and tocotrienols

Jamila M. Al-Saqer^a, Jiwan S. Sidhu^{a,*}, Suad N. Al-Hooti^a, Hanan A. Al-Amiri^a, Amani Al-Othman^b, Latifa Al-Haji^c, Nissar Ahmed^c, Isa B. Mansour^d, Johari Minal^d

*Berechning: Department, Kavail Institute for Scientife Research PO Bas 24855, 11199 Safer, Kavail *Systems Development Department, Kavail Institute for Scientife Research PO Bas 24855, 11199 Safer, Kavail *Centrel Analysis Alachesteries, Kavail Institute for Scientife Research, PO Bas 2585, 11199 Safer, Kavail *Alalysium Polin Gd Based, PO Bas 18650, 50750 Knile Lampur, Maleysia

Received 1 November 2002, received in revised form 24 April 2003; accepted 14 August 200

Abstract

Two functional floods (pan bread and sugar-map cookies), were proputed using red paths orbits (BFOA) and red plans shortening (BFOS) with the objective of providing higher amounts of antisolatal vasuame E now office. The bread statum E control is the polin observation in the polin obs and in this study were significantly higher (traping from 71.78 to 11.75 mg kg fat) than those in the polin observations for inpuls observed the policy of the polic

JONG EINCHET LIG. An rights reserved.
 Kryworth Functional Josebs, Red pales ofeirs, Vitamin E; Tocopherolis, Tocopherolis, Braud, Cookies; Notritive val.



What functional food is the ideal for this ingredient?

CONCLUSION: Functional food ingredient?

- Any food with lipids
 - •Especially, food with PUFA
 - •New enriched omega-3 food
- •Combination with other antioxidants
 - Synergistic effects
- •No regulation problem, natural and frequent ingredient
- •Important to combine with other active components in new functional foods

TOCOPHEROLS



CONCLUSION:

VITAMIN E...

...Functional food ingredient?

FUNCTIONAL INGREDIENTS

Lipidic Ingredients from vegetable products

Tocopherols

Sterols Carotenoids

Other antioxidants

What are they? Structure - types-function

Why functional ingredients? (bioactivity)

Do you have scientific evidence? Articles

Do you have clinical studies? Health claims



OK, Lets do it

Where are they? Sources-industrial production

What functional food is the ideal for this ingredient? Available, compatible, complementary

Produce and sell it!

STEROLS



PHYTOSTEROLS.....

STEROLS STANOLS STEROL ESTERS **STANOL ESTERS**

...added EVERYWHERE






What are they? Structure - classes

- Sterols and Stanols
- Sterol esters and stanol Esters
- Minor components in oils (unsaponifiable)
- Both tocopherols and carotenoids are groups of well known compounds with interesting activity
- Very common, frequent in small amounts
- Not clear beneficial activity



STEROLS





β-Sitosterol



• Structure: Sterols and Stanols

•Phytosterols: Sterol and stanol Esters

•Main activity: Inhibit cholesterol absorption

•Compete in micelles for absorption of lipids with cholesterol

·Natural, mostly sterols

•STANOL: no absorption

•Diverse family - similar behaviour

Industrial sources:

•tall oil

deodorizer oil distillates

Hydrogenated to produce stanols

STEROLS What are they? Structure - types-function

• Structure: Sterols and Stanols

FIGURE 1. Examples of basic structures of free sterol and steryl conjugates. R (side chain) varies between different sterols.


glycoside



Sterols and Stanols

D-5-avenasterol


beta-sitosterol



STEROLS

Why functional ingredients? (bioactivity)

- Main biological effect: competition in micelles
- Other demonstrated effects: LDL-c





- · Phytosterols:
 - Sterols and Stanols
 - Sterol Esters and stanol Esters
- •Main activity:
- •Reduce the absorption of cholesterol from the gut and so lower serum concentrations of cholesterol
- •Plant sterols or stanols that have been esterified to increase their lipid solubility can be incorporated into foods




STEROLS

Atherosclerosis

- hardening of the arteries due to the deposition of atheromas
- heart disease is the leading cause of death
- caused by the deposition of cholesteryl esters on the walls of arteries
- atherosclerosis is correlated with high LDL and low HDL







- •Sterols Hydrogenated to produce stanols
- Free Sterols Esterified to produce sterol esters (stanol esters)

Sterol Esters and stanol Esters

- •Plant sterols or stanols that have been esterified to increase their lipid solubility can be incorporated into foods
- •Diverse family similar behaviour

Main activity:

Reduce the absorption of cholesterol from the gut and so lower serum concentrations of cholesterol

STEROLS

What are they? Structure - types-function

FIGURE 1. Examples of basic structures of free sterol and steryl conjugates. R (side chain) varies between different sterols.





- •Chemical characterization and Different tests (activity *in-vitro*...): many
- •Epidemiological studies: several evidences
- •Animal model tests (in-vivo)
- Pre-clinical: Dose effects, toxicological risks, etc.: studied
- •Clinical studies, Intervention studies: many

Stanols and sterols have been well tolerated in numerous clinical trials on over 1800 people, with doses up to 25 g per day.

Studies have established that at least 1 gram of stanol and sterol in ester form can result in a statistically significant reduction in cholesterol.

Recent studies of plant sterol and stanol esters in humans have shown, however, that maximum cholesterol-lowering benefits are achieved at doses of 2-3 g per day.



STEROLS

Is there any scientific evidence? Articles

- •Chemical characterization and Different tests (activity *invitro...*): many
- •Epidemiological studies: several evidences
- •Animal model tests (in-vivo)
- Pre-clinical: Dose effects, toxicological risks, etc.: studied
- •Clinical studies, Intervention studies: many
- •Reliability: Total
- •Controversial: very little for a new product
 - None on effect
 - •Some on high dose supplementation effect
 - •Much on interaction with fat-soluble vitamins and carotenoids
- Accepted: completely
- •Approved health claims: yes, Ok (US, UE, etc.)



Where are they? Sources

- · Phytosterols:
 - Sterols and Stanols
 - Sterol Esters and stanol Esters
- Industrial sources:
 - •tall oil
 - deodorizer oil distillates
- Hydrogenated to produce stanols
- Sterol Esters and stanol Esters
- •Plant sterols or stanols that have been esterified to increase their lipid solubility can be incorporated into foods



STEROLS

Are ther clinical studies? Health claims

Properties: (When at least 1 gram per day is consumed in ester form for statistically significant effects)

- Reduce total cholesterol
- Reduce LDL cholesterol
- •Do not affect HDL cholesterol or triglycerides



"Foods that may qualify for the health claim based on plant sterol ester content include spreads and salad dressings. Among the foods that may qualify for claims based on plant stanol ester content are spreads, salad dressings, snack bars, and dietary supplements in softgel form. Foods that carry the claim must also meet the requirements for low saturated fat and low cholesterol, and must also contain no more than 13 grams of total fat per serving and per 50 grams." (FDA, 2000)



Health claims:

Diets low in saturated fat and cholesterol that include at least 1.3 grams of plant sterol esters or 3.4 grams of plant stanol esters, consumed in 2 meals with other foods, may reduce the risk of heart disease. In addition to and in the presence of the health claim, products with plant stanol/sterol esters may also state that they can reduce cholesterol levels.



Plant sterols first came on the European market in 1995 in a fortified margarine called Benecol, launched in Finland by Raisio Life Sciences. But even after hitting US shores in 1999 by McNeil (Benecol) and Unilever (Take Control), sales did not live up to expectations. The cholesterol-lowering potential of stanols and sterols had been known for 50 years, and with an estimated 105 million Americans with borderline or high cholesterol, the market potential certainly was there. What happened?

The trouble was, margarines are not commonly associated in consumers' minds with healthful eating, and in America at least, spreads are not routinely found on the daily dinner table.

In 2002, the Food & Drug Administration finally allowed the cholesterol-lowering heart-health claim for plant sterols to be available to a wider range of foods, and if the flurry of product launches of the past few years is any indication, the train has finally taken off.



STEROLS

Are ther clinical studies? Health claims

- Accepted health claims: Now in a second phase:
- LONG TERM STUDIES
- Possible problems for very large daily ingestion
- Interactions:

M. Richelle, M. Enslen, C. Hager, M. Groux, I. Tavazzi, J.-P. Godin, A. Berger, S. Metairon, S. Quaile, C. Piguet-Welsch, L. Sagalowicz, H. Green, and L. B. Fay Both free and esterified plant sterols reduce cholesterol absorption and the bioavailability of {beta}-carotene and {alpha}-tocopherol in normocholesterolemic humans

Am. J. Clinical Nutrition, July 1, 2004; 80(1): 171 - 177.

Noakes, M., Clifton, P., Ntanios, F., Shrapnel, W., Record, I. & McInerney, J. (2002) An increase in dietary carotenoids when consuming plant sterols or stanols is effective in maintaining plasma carotenoid concentrations. Am. J. Clin. Nutr. p.79



PHYTOSTEROLS.....

STEROLS STANOLS STEROL ESTERS STANOL ESTERS

...added EVERYWHERE



Lipidic Ingredients from vegetable products

Tocopherols

Sterols

Carotenoids

What are they? Structure - types-function

Why functional ingredients? (bioactivity)

Do you have scientific evidence? Articles

Do you have clinical studies? Health claims



OK, Lets do it

Where are they? Sources-industrial production

What functional food is the ideal for this ingredient? Available, compatible, complementary

Produce and sell it!

FUNCTIONAL INGREDIENTS



Tocoferoles Carotenoides Esteroles


Complexity: Sterols (like one)-Tocos (4+4)-Carotenoids (100s)

Level of scientific evidence? Articles-controversy

Status of clinical studies? Health claims posed

Sources-industrial production:
Tall oil (sterols) DOD (sterols+toco) Plants and algae (carotenoids)

What functional food is the ideal for this ingredient?
Available, compatible, complementary
Different cases:tocos-sterols-carots-etc







• Structure

CAROTENOIDS



What are they? Structure - types-function

• Structure



Where are they? Sources-industrial production

Vegetable products



•Natural Mixed Carotenoids family of naturally occurring yellow, orange and red pigments found in various fruits, cruciferous, yellow and dark -

found in various fruits, cruciferous, yellow and dark - green vegetables and also abundantly in algae.

What are they? Structure - types-function • Structure | Canthexanihia | Cant






Structure

CAROTENOIDS

What are they? Structure - types-function

• Structure: Carotenes and xanthophylls








Activity: Carotenes and xanthophylls

- •Main activity: antioxidant
- •Pro Vitamin A
- •Natural (some synthetic)
- •Diverse and complex family





CAROTENOIDS



Why functional ingredients? (bioactivity)

- Main biological effect:
- Other demonstrated effect:
- Additional physiologic effects beyond vitamin A
 - Serving as an "antioxidant"
 - Remove excess "electrons" from cell system
 - Electrons (free radicals) damage cells and DNA
 - Can cause mutations
 - Protecting from cancer (related to antioxidant function?)
 - Protecting from heart disease?
 - Supports immune system function



CAROTENOIDS

Carotenoids. Roles in the Body

- Promote vision (retinal)
 - Maintenance of the cornea
 - Converting light energy into nerve impulses
- Participate in protein synthesis and cell differentiation
 - Differentiation of epithelial and goblet cells (retinoic acid)
- Support reproduction and growth
- Support immunity (retinoic acid and carotenoids)
- · Involved in bone growth and remodeling
- Synthesis of glycoproteins
- Antioxidant activity (B-carotene....)





Do you have clinical studies? Health claims

- Pre-clinical for some of them
- •Different tests (in-vitro, in-vivo...)
 - Agreement on antioxidant activity
- Epidemiological studies form natural mixtures of carotenoids
- Clinical studies:
- Controversial: Strong
 - •Problem with beta-carotene and smokers
- •Enrichment Accepted: ??
- Accepted health claims:



CAROTENOIDS

Where are they? Sources

Sources: Carotenes and xanthophylls

- •Natural (some synthetic)
- Diverse and complex family
- Industrial sources:



biotechnological production with yeasts/fermentation microalgae (Dunaliella, Haematoccocus, etc.) chemically synthesized





Different cases:tocos-sterols-carots-etc

Tocoferoles Carotenoides

Esteroles

Complexity: Sterols (like one)-Tocos (4+4)-Carotenoids (100s)

Level of scientific evidence? Articles-controversy

Status of clinical studies? Health claims posed

Sources-industrial production:
Tall oil (sterols) DOD (sterols+toco) Plants and algae (carotenoids)

What functional food is the ideal for this ingredient?
Available, compatible, complementary
Different cases:tocos-sterols-carots-etc

CAROTENOIDS

Lycopene from tomato

Structure of lycopene



Lycopene

CAROTENOIDS



Lycopene from tomato

- · Biological activity
 - Antioxidant
 - Efficient oxygen quencher and trap peroxyl radicals
- •Fresh tomatoes:
 - •80-90% of total carotenoids
 - all-trans-lycopene
 - •0,7-20 mg/100g Wet basis



Processed tomatoes: increased cis-lycopene (4-27%)

- •Human serum: > 50% cis-lycopene
- •Skin and pericarp rich in lycopene ---industrial waste material (Al-Wandawi, 1985)

CAROTENOIDS



Lycopene from tomato

- Biological activity
 - Antioxidant
 - Protection of DN...
- - fikoobane
- Lycopene content affected by agricultural practices:
 - •irrigation, fertilization, temperature and light, variety, degree of maturity at harvest, harvest time, postharvest storage (Shi, 2002)
- ·Bioavaliability for humans:

Isomeric form food matrix lipid content other interactions (fiber)



CAROTENOIDS



Lycopene from tomato

- Biological activity
 - Antioxidant

- Health promoting effect
 - Protection against some cancers:
 - Prostate, breast, digestive tract, epithelial... (Colditz, 1985, Gerster, 1997)
 - •Studies (90's) on tomato consumption and protection against digestive cancer (Franchesi, 1994,
 - •Giovannucci, 1995, prostate cancer in 48,000 men / 4 years / 46 fruits&vegetables
 - •lower risk strongly associated with tomato sauce comsumption
 - Atherosclerosis and Coronary Heart Disease
 - •Reduce LDL oxidation (Kristenson, 1997)
- •Recomendación diaria 10-50 mg

CAROTENOIDS

Lycopene from tomato



Roche Lycopene Self-Affirmed GRAS

PARSIPPANY, N.J.-Roche Vitamins Inc. announced it has self-affirmed its ! GRAS (Generally Recognized as Safe) through an independent panel evaluation been scientifically linked to reducing the risk of certain cancers, as well as impro lung, cardiovascular and stomach health, according to the company. Roche lyco in both water-dispersible and oil-soluble forms and is suited for an array of proc nutritional bars, beverages and dairy products. Roche (<u>www.roche.com</u>) noted stable, easy to process and will not affect the taste profile of food and beverage





<u>Mission Statement</u>
LycoRed Natural Products Industries Ltd., is a research and development company dedicated to the scientific and commercial development, production and and commercial overlopment, production and marketing of natural products based on the phytonutrients from tomato, including the carotenoid lycopene, beta-carotene, phytoene and phytofluene, phytosterols and other important phytochemicals.



About the Company
LycoRed is a subsidiary of Makhteshim Agan Industries,
Ltd., one of Israel's largest industrial groups, a public
company, with a turnover in 2001 of more than \$500
million. LycoRed supervises production of

LYC-O-MATO® tomato extract from the field to the LVC-0-MATO* tomote obtact from the field to the finished product. Coordination of the commercial growing of a non-genetically modified, bycopen-wich variety of tomato plant is just the start of the multi-stepped process that assures high variety quality. Phytichemicals are extracted using proprietary processing sethnology. Specialised extracted using proprietary pracking and storing behinques assure a long shelf life. LycoRed markets LYC-O-MATO[®], its scientifically standardized, natural tomato extract, worldwide.



LycoRed's LYC-O-MATO® form functional properties supported by basic research and clinical research. A growing body of scientific evidence



Lycopene from tomato

- · Extraction of lycopene from tomatoes
- •Need of high purity lycopene products -·food safety regulations
- Need of further studies on toxicology and bioavailability
- ·Large amount of skin and pericarp discarded waste of peeling



- Processed tomatoes: increased cis-lycopene (4-27%)
- •New extraction technologies under development
 - complex bioactive components
 - •sensitive to oxygen, thermal, (oxidation, isomerization, etc.)

CAROTENOIDS




Lycopene from tomato waste

Limitation: reliable Industrial source Difficult extraction

2638 J. Agric. Food Chem. 2002, 50, 2638-2643





N. L. Rozzi, $^{\uparrow}$ R. K. Singh, $^{\sharp}$ R. A. Vierling, § and B. A. Watkins $^{\bullet, \uparrow}$

Tomato sects and skins acquired from the hypercluit of a local tomato processing facility were studied for supervirtical fluid extraction (SEE) of hyptochemicals. The extracts were analyzed for hypopen-client processing facility were fluid for supervirtical fluid extraction (SEE) of hyptochemicals. The extracts were analyzed for hypopen-client fluid chromatography-electrochemical detection and compresed to a chemically extracted control. SEE were certained using CO; at severe temperatures (24.78—68.26 MPs). The effect of CO; See wate and volume also was investigated. The results indicated that he percentage of program extracted chromated with elevated temperature and pressure until a sextended decreased. Conditions for the optimum extraction of hypopens from 3 g of raw material were determined to be 65 CO, 347 MPs, and 550 mL; of CO; at flow mit exit of 55 mL/mls. These conditions resulted in the extraction of 61.0% of the lycopene (7.19 μg lycopeneig).



percritical fluid CO₂ has been used to extract α- and rotten and lycopene from several plant sources (14-17), sili and Callahan (14) compared supercritical fluid extraction



Other natural sources

- •Not synthetic—not single carotenoid:
- Natural Mixed Carotenoids
- •Carrots, corn, colored vegetables....
- •No industrial scale extraction



- •Microalgae are a new source of carotenoids and other bioactive ingredients
 - Chlorella,
 - Spirulina,
 - Dunaliella,
 - · Microalgae Haematococcus pluvialis,
 - etc.



CAROTENOIDS Other natural sources WHY TO USE ALGAE?

Man has been using algae since long time ago (thousand years) with different purposes.

In China they have been used since 2700 a.c.; greeks and romans used algae as food, as medicinal plants and cosmetics.

Aztecs used Spirulina, harvested from Texcoco Lake, as proteic supplement.


In the world, around 7 billion Tons of algae (net weigth) are processed, being asian countries the main producers (around 80% of the raw material).



Other natural sources

WHY TO STUDY MICROALGAE?





The photosynthetic activity of the algae generates O2, to avoid attack of O2 the microalgae accumulate <u>antioxidant compounds</u>


Some compounds with <u>antiviral</u> and <u>antimicrobial</u> activity have been described

- At present, research is conducted to verify the efficiency of algae against certain diseases and illness: trombosis, cholesterol, rheumatism
- The main consideration is safety-toxicity, any possibility of toxicity precludes the use of the microalgae.
- The literature shows that ß-carotene and PUFAS have been widely studies, while other compounds with potential activities (antioxidant, antimicrobial or antiviral (polyphenols, peptides, polysaccharides)) have not been studied yet.
- Commercial availability

CAROTENOIDS

Astaxanthin from microalgae

- •Microalgae are a new source of carotenoids and other bioactive ingredients Chlorella, Spirulina, Dunaliella, etc.
- Biological activity
 - Antioxidant
 - Efficient oxygen quencher
 - trap peroxyl radicals



- Microalgae Haematococcus pluvialis:









Extraction of astaxanthin from Haematococcus

Antioxidant more powerful than other carotenoids Benefits for carciovascular health for macular degeneration (eye health)



Haematoccocus pluvialis

(Empresas: US Nutra (LaHaye), Separex, Fuji, Cyanotech...)








"As an antioxidant, astaxanthin has been shown to be up to 500 times more powerful than Vitamin E and more than 10 times as potent as beta-carotene."









Beta-carotene from microalgae: Dunaliella

- •Microalgae are a new source of carotenoids and other bioactive ingredients Chlorella, Spirulina, Dunaliella, etc.
- Biological activity
 - Antioxidant
 - Efficient oxygen quencher
 - trap peroxyl radicals





•Microalgae Dunaliella salina


Is a single celled, salt-water micro-algae accumulates massive amounts of carotenoids under appropriate growth conditions.

Natural Mixed Carotenoids found in Dunaliella salina are among nature's best antioxidants, containing a variety of carotenoids including β -Carotene, α -carotene and Xanthophylls like Zeaxanthin, Cryptoxanthin and Lutein.

CAROTENOIDS







- ullet Composition related to different activity of eta-carotene as antioxidant
- Higher antioxidant activity of 9-cis isomer in vitro (Levin and Mokady, Free Radical Biology and Medicine, 17, 77-82 (1994))
- Better activity of <code>Dunaliella</code> extracts vs synthetic β -carotene in protecting hepathic diseases in vivo



Functional Compounds from **Dunaliella salina**:(SFE, ASE)

- High ratios 9 cis/ all trans β -carotene isomers could be obtained tuning the SFE conditions- studies about the relationship with antioxidant activity are being conducted in our laboratory via in vitro assays
- High yield were obtained by SFE of *Dunaliella salina* microalgae (5-10% for CO2 or CO2 + oil)
- Antimicrobial activity was obtained for PLE extracts of *Dunaliella salina*
- PLE of Dunaliella with ethanol provide good antimicrobial extracts using a green solvent
- The compounds associated to the antimicrobial activity were mainly fatty acids and ionones coming from carotenoid decomposition





ANTIOXIDANTS



Antioxidants from Spirulina platensis (SFE, SWE)



- Blue-Green microalga belonging Cyanobacterium
- Non toxic / Edible
- Live in high salinity lakes
- Up to 60 -70 % dry weight is protein
- Phycocyanins are the main proteins: several functional activities.
- Relatively high content GLA
- Chlorophyll a, xantophylls and other carotenoids are their main photosynthetic pigments

















What are they? Structure - types-function

· Antioxidant capacity of carotenoids



J. Agric. Food Chem. 2002, 50, 221-226 221


Trolox Equivalent Antioxidant Capacity of Different Geometrical Isomers of α -Carotene, β -Carotene, Lycopene, and Zeaxanthin

Volker Bohm, * , $^{\uparrow,\downarrow}$ Ni Luh Puspitasari-Nienaber, $^{\downarrow}$ Mario G. Ferruzzi, $^{\downarrow}$ and Steven J. Schwartz $^{\downarrow}$

Institute of Nutrition, Friedrich Schiller University Jena, Dornburger Strasse 25-29, 07743 Jena, Germany, and Department of Food Science and Technology, The Ohio State University, 2001 Fyffe Court, Columbus, Ohio 43210-1097

Isomerization of carotenoids, which is often encountered in food processing under the influence of temperature and light, may play a role in the observed protective effects of this group of secondary plant products. Investigation of in vitro antioxidant activity of prominent carotenoid geometrical isomers was undertaken in light of recent reports Illustrating a large percentage of carotenoid (2)-isomers in biological fluids and tissues. a Crarotene, (3-carotene, lycopene, and zeaxanthin were isolated from foods or supplements and subsequently photoisomerized with iodine as a catalyst. Major 2-isomers of each carofenoid were fractionated by semicrogramtive C.- HPI C. In vitra militoxiant activity of all To date, published results for the antioxidant activity of β -carotene isomers differ widely due to the various test systems employed.

CAROTENOIDS What are they? Structure - types-function Antioxidant balance Vitamin C Uric acid Bilirubin Blood Ceruloplasmin (binds copper) Transferrin (binds iron) Peroxisome Cytosol of Zinc-copper-Catalase Vitamin C body cell superoxide dismutase Manganesesuperoxide dismutase Glutathione peroxidase Nucleus (selenium) DNA repair Vitamin E Carotenoids enzymes








What are they? Structure - types-function

Antioxidants in food





ANTIOXIDANTS

What are they? Structure - types-function

Free Radical Damage

- Cancer: prostate, cervical, breast, lung
- · Heart disease
- · Immune system degeneration
- Stroke, cerebrovascular disease
- · Respiratory function
- · Cataracts, age-related macular degeneration
- · Memory and cognitive function
- · Sun-damaged skin, wrinkles
- · Diabetes complications



Oxidative stress




What are they? Structure - types-function

Free Radical Damage

Where Do Free Radicals Come From?

They form and accumulate in our bodies as a result of modern life:

- Cigarette smoke
- Air pollutants, such as ozone, nitrogen dioxide
- Some food preservatives, including nitrites
- Aerobic activity
- Metabolism of fats





Antioxidant potency of Rosemary diterpenes

- •Phenolic diterpenes: only present in Rosemary (Rosmarinus officinalis) and Sage (Salvia officinalis)
 - •Native from Mediterranean area and South America
- •Main component in Rosemary and Salvia: carnosic acid
- Antiox. activity depends on:
 - •season, climatic conditions, light, harvesting, storage, etc.
 - •Extraction: best results with supercritical CO2



Antioxidant potency of Rosemary diterpenes carnosic acid rosmarinic acid rosmadial Cuvelier et al., 1996)



Why functional ingredients? (bioactivity)


Primary antioxidants

$$ROO + AH = ROOH + A$$



ANTIOXIDANTS

Antioxidant potency of Rosemary diterpenes



- •Extraction of carnosic acid with supercritical CO2
- •Main component in Rosemary and Salvia
- Commercial extracts
- •Clinical studies: Not yet
- •Accepted health claims: Not yet
- •Functional food: in market





Antioxidant potency of Rosemary diterpenes

- •Rosemary (Rosmarinus officinalis) Phenolic diterpenes
- •Extraction of carnosic acid with supercritical CO2

European Food Research and Technology

DOI 10.1007/s00217-005-1139-v

Original Paper

In vitro antioxidant analysis of supercritical fluid extracts from rosemary (Rosmarinus officinalis L.)

Sofía Cavero · Laura Jaime · Pedro J. Martín-Álvarez · F. Javier Señoráns · Guillermo Reglero · Elena Ibañez (♥)

S. Cavero · L. Jaime · F. J. Señoráns · G. Reglero

Área de Tecnología de Alimentos, Facultad de Ciencias, Universidad Autónoma de Madrid, 28049 Cantoblanco, Madrid, Spain

P. J. Martín-Álvarez · E. Ibañez

Departamento de Caracterización de Alimentos, Instituto de Fermentaciones Industriales (CSIC), Juan de la Cierva 3, 28006 Madrid, Spain





ANTIOXIDANTS

Antioxidant potency of Rosemary diterpenes

- Rosemary (Rosmarinus officinalis) Phenolic diterpenes
- •Extraction of carnosic acid with subcritical water

AGRICULTURAL AND FOOD CHEMISTRY

J. Agnic. Food Chem. 2003, 51, 375-382 375

Subcritical Water Extraction of Antioxidant Compounds from Rosemary Plants

ELENA IBAÑEZ,*:† ALENA KUBÁTOVÁ,‡ F. JAVIER SEÑORÁNS,§ SOFIA CAVERO,§

Instituto de Fermantaciones Industriales, CSIC, Juan de la Cierva 3, 28006 Madrid, Spain, Energy and Environmental Research, Center, University of North Daloto, Campus Bers 9018, Grand Ferk, North Daloto 58202, and Avas de Tocnologia de Alimantoc, Universidad Antonoma de Madrid, Cinded Universitatia de Cantolhanco, 20049 Madrid, Spain

Subcritical water entraction at several temperatures ranging from 25 to 200 °C has been studied to selectively extract anticidation compounds from rosemary leaves. An enhancise characterization of the fractions obtained using subcritical water at different temperatures has been carried out by LC-MS, and the anticidant activities of the extracts have been measured by a five radical method (OPPH). MS, and the anticidant activities of the extracts have been measured by a five radical method (OPPH). Results indicate byte selectivity of the subcritical vater toward the most active compounds of rosemary such as camosol, rosemand, camosol acid, methyl camosate, and some fishronoids such as originatific and genkvanin. The anticidical ractivity of the fractions obtained by extraction at different water temperatures was very high, with values around 11.3 µg/mL, comparable to those achieved by SFE of rosemary leaves. A study of the effect of the temperature of the extraction efficiency of the most typical rosemary anticioidant compounds has been performed.

KEYWORDS: Antioxidant activity; antioxidant compounds; rosemary plants; SFE; subcritical water extraction





FUNCTIONAL INGREDIENTS



FUNCTIONAL FOODS

Lipophilic Ingredients from vegetable products

Bioactive and health promoting ingredients from vegetable products



Potential functional food ingredients





Functional Ingredients from vegetable products

Bioactive and health promoting ingredients from vegetable products

- 1. FATTY ACIDS (OLEIC, LINOLENIC, OMEGA-3, ETC.)
- 2. TOCOPHEROLS
- 3. STEROLS
- 4. CAROTENOIDS
- 5. OTHER ANTIOXIDANTS: PHENOLIC DITERPENES
- 6. ISOFLAVONES AND ISOFLAVONOIDS
- 7. FLAVONOIDS
- 8. PHENOLIC COMPOUNDS
- 9. ORGANOSULFUR COMPOUNDS
- 10. ETC...

FUNCTIONAL INGREDIENTS



Other Candidates:

FUNCTIONAL INGREDIENTS



OPPORTUNITIES

Where to find: Other possibilities

Look around:

vegetables fruits

non-usual edible oils

spices

Think different microalgae mushrooms berries

...??







FUNCTIONAL INGREDIENTS

Other candidates



There is no question that consumer awareness for antioxidants is high, but knowledge gaps still exist, especially with regard to testing.

By Rebecca Wright Editor

FUNCTIONAL INGREDIENTS

OPPORTUNITIES

New ingredients Critical points

Safety

Evidence

Availability

Regulations

Extraction of compour _______

Market









FUNCTIONAL INGREDIENTS

Any case is Different: tocos-sterols-carots-etc

How to start? Local product (food) Local source of ingredients (Rosemary case)

Characterize source and components. Isolate (extract-fractionate) active compounds

Test them. Chemical-invitro-biological activityin vivo-clinical: scientific evidence? Articles

Status of clinical studies? Health claims

Source availabilitypossibility of industrial production:

What functional food is the ideal for this ingredient? Available, compatible, complementary Produce and sell it!



Research group in functional foods Food Science Department University Autónoma of Madrid (UAM)



Contact: F. Javier Señoráns javier.senorans@uam.es

http://www.uam.es/javier.senorans

Sec. Dept. de Ciencias de la Alimentación, UAM Facultad de Ciencias, C-XVI, 502 28049 Madrid (Spain)

http://www.uam.es/otros/gruta/Research/Principal.htm

www.uam.es/tecnol-alim

